



## 80 mg Spot-on Solution for Large Cats & Pet Rabbits

### Description:

Each tube contains 0.8 ml of a 10 % w/v cutaneous solution of imidacloprid (equivalent to 80 mg imidacloprid/tube).

### Indication(s)

For the prevention and treatment of fleas on cats of 4 kg and over and treatment of fleas on pet rabbits 4 kg and over

### Contraindications

Do not use on rabbits intended for human consumption. Do not use this product on cats or pet rabbits weighing less than 4 kg.

Do not treat unweaned kittens of less than 8 weeks of age or rabbits of less than 10 weeks of age. Do not allow recently treated animals to groom each other. Care should be taken to avoid the contents of the tube coming into contact with eyes or mouth of recipient animal. Do not use in animals that are known to be hypersensitive to the active substance or any of the excipients.

### Target Species:




Cats and Rabbits

### Dosage and administration:

One tube should be applied to cats and pet rabbits weighing 4 kg and over (equivalent to a maximum dose of 20 mg/kg imidacloprid). This dose will provide protection against flea infestation in the cat for 3–4 weeks and up to 1 week on the rabbit.


The product can be used as part of a treatment strategy for the control of Flea Allergy Dermatitis (FAD) in cats where this has been previously diagnosed by a veterinary surgeon.

Hold tube in an upright position, twist and pull off the cap. Remove one tube from the package. Use reversed cap to twist and remove seal from tube.



### Administration

Part the hair on the pet's neck at the base of the skull until the skin is visible.



Place the tip of the tube on the skin and squeeze firmly several times to empty its contents directly onto the skin. Do not rub in. Apply only to undamaged skin. For optimal prevention and control of flea infestation, monthly treatments are recommended. The efficacy of the product may be reduced if the animal becomes markedly wet. If retreatment becomes necessary earlier than 4 weeks, treatment may be repeated. Do not apply more frequently than weekly. The product is bitter-tasting and salivation may occasionally occur if the cat licks the application site immediately after treatment. This is not a sign of intoxication and disappears within a few minutes without treatment. Correct application at the base of the skull will minimise the opportunity for the pet to lick the product

### Withdrawal period

Not Applicable

### Special storage precautions

KEEP OUT OF REACH OF CHILDREN

Store away from food, drink or animal feeding stuffs. Do not use after expiry date. After use, replace cap on tube.

### Special warning(s)

FOR EXTERNAL USE ONLY  
FOR ANIMAL TREATMENT ONLY

**Special precautions to be taken by the person administering the veterinary medicinal product to animals:**

Do not eat, drink or smoke during application. Wash hands thoroughly after use. Wash off any skin contamination with soap and water. People with known skin sensitivity may be particularly sensitive to the product. This product contains benzyl alcohol and may cause skin sensitisation or transient skin reactions (e.g. allergy, irritation, tingling). Avoid contact between the product and the eyes and mouth. If the product gets into eyes accidentally, the eyes should be thoroughly flushed with water. If skin or eye irritation persist, obtain medical attention. If the product is accidentally swallowed, obtain medical attention immediately. After application do not stroke or groom animals until the application site is dry (typically within an hour or so).

**General Precautions:**

Re-infestation from emergence of new fleas in the environment may continue for 6 weeks or longer after treatment. To aid reduction in environmental challenge the additional use of a suitable environmental treatment against adult fleas and their developing stages is recommended. Poisoning following inadvertent oral uptake in animals is unlikely. In this event, treatment should be symptomatic under veterinary medical attention. There is no known specific antidote but administration of activated charcoal may be beneficial. The solvent in Advantage® may mark certain materials including leather, fabrics, plastics and finished surfaces. Allow application site to dry before permitting contact with such materials.

**Pregnancy and Lactation:**

No primary or embryotoxic, teratogenic or reproductive toxic effects have been observed during the studies with imidacloprid on rats and rabbits. Studies on pregnant and lactating queens together with their offspring are limited. Evidence so far suggests that no adverse effects are to be expected in these animals.

**Interaction with other medicinal products:**

No incompatibility has been observed between this product at twice the recommended dose and the following commonly used veterinary products: lufenuron, pyrantel and praziquantel (cats). The compatibility of the product was also demonstrated with a wide range of routine treatments under field conditions including vaccination.

**Special precautions for the disposal of unused product or waste materials, if any:**

Dispose of used packaging in the household refuse. Unused product should be returned to the veterinary surgeon.

**Other Information**

Packaging style Blister packs containing either 2, 3, 4 or 6 unit dose tubes or a single unit dose tube without blister.

Pack Size Carton contains 1 tube or 1, 5, 10 or 20 blisters each with 2, 3, 4 or 6 tubes

Not all pack sizes may be marketed.

In recent studies, in addition to the adulticide flea efficacy of imidacloprid, a larvicidal flea efficacy in the surroundings of the treated pet has been demonstrated. Larval stages in the pet's surroundings are killed following contact with a treated animal.

**Name and address of the Marketing Authorisation Holder and of the Manufacturer:**

NFA-VPS

**Marketing Authorisation:**

The Marketing Authorisation number is Vm 00010/4134

**Marketing Authorisation Holder:**

Bayer plc, 400 South Oak Way, Green Park, Reading, RG2 6AD,  
Tel: 0118 206 3000

**Manufactured by:**

KVP Pharma + Veterinär Produkte GmbH, Projensdorfer Str. 324,  
D-24106 Kiel, Germany

**The name of this veterinary medicine is:**

Advantage 80 mg Spot-on Solution for Large  
Cats & Pet Rabbits

® Registered Trademark of Bayer AG

Date of Leaflet Preparation: April 2019

